



STUK-B 286 / JUNE 2022

Virtanen S, Vartti V-P, Turunen J, Huusela K, Teräväinen M,
Torvela T, Mattila A

B



Monitoring of radioactivity in the environment of Finnish nuclear power plants

Annual report 2021

Radiation and Nuclear Safety Authority
Jokiniemenkuja 1
FI-01370 Vantaa, Finland
www.stuk.fi

Further information:
Sinikka Virtanen
sinikka.virtanen@stuk.fi
telephone +358 9 759 88 556

ISBN 978-952-309-533-5 (pdf)
ISSN 2243-1896

VIRTANEN Sinikka, VARTTI Vesa-Pekka, TURUNEN Jani, HUUSELA Kari, TERÄVÄINEN Mikko, TORVELA Tiina, MATTILA Aleks, Monitoring of radioactivity in the environment of Finnish nuclear power plants. Annual report 2021. STUK-B 286 Helsinki, 2022, 49 p.

KEY WORDS: Environmental monitoring, nuclear facility



Contents

1 SUMMARY	5
2 INTRODUCTION	5
3 RELEASES FROM NUCLEAR POWER PLANTS	6
4 ENVIRONMENTAL MONITORING PROGRAMME OF THE LICENSEE	7
5 ENVIRONMENTAL MONITORING PROGRAMME AND METHODS OF THE RADIATION AND NUCLEAR SAFETY AUTHORITY	8
5.1 MONITORED PATHWAYS AND SAMPLING	9
5.1.1 OUTDOOR AIR AND ATMOSPHERIC DEPOSITION	9
5.1.2 TERRESTRIAL ENVIRONMENT	10
5.1.3 MARINE ENVIRONMENT	11
5.1.4 INHABITANTS OF THE SURROUNDINGS	15
5.2 MONITORING METHODS	16
5.2.1 SAMPLE PROCESSING AND ANALYSIS	16
6 RESULTS OF ENVIRONMENTAL MONITORING	19
6.1 OUTDOOR AIR AND ATMOSPHERIC DEPOSITION	19
6.2 TERRESTRIAL ENVIRONMENT	20
6.3 MARINE ENVIRONMENT	27
6.4 INHABITANTS OF THE SURROUNDINGS	39
7 SUMMARY AND CONCLUSIONS	40
8 REFERENCES	41
9 ANNEXES	41

1 Summary

This report describes the results of radiation monitoring carried out by the Radiation and Nuclear Safety Authority (STUK) in the environment of the Loviisa and Olkiluoto nuclear power plants in 2021. STUK's environmental monitoring and measurement activities complement and verify the environmental monitoring and release measurements conducted by the power plants. The monitoring is implemented by collecting samples from the terrestrial and marine environment in the vicinity of the power plants and of outdoor air. In addition, the concentrations of radioactive substances in the bodies of inhabitants of the surrounding area of the power plant are monitored. The environmental samples are analysed in STUK's laboratory. The radioactive substances contained in the collected samples are determined by gamma spectrometric and radiochemical analysis methods.

In some of the collected samples, small quantities of radioactive substances originating from the power plant were found. There was no significant deviation from the environmental findings of the previous years in terms of the identified radioactive substances or their quantities. Radioactivity originating from the power plant observed in the environment is insignificant in terms of radiation exposure of the environment and people. The results of the release measurements reported by the nuclear power plants and the findings of the environmental monitoring carried out by the nuclear power plants are compatible with the findings made by STUK as part of its own monitoring.

2 Introduction

The use of nuclear energy is prescribed for in the Nuclear Energy Act (990/1987) and Nuclear Energy Decree (161/1988). Under Section 7 c(1) of the Nuclear Energy Act, *releases of radioactive substances caused by the use of nuclear energy shall be restricted in compliance with the optimisation principle of radiation protection laid down in Section 6 of the Radiation Act (859/2018). In the optimisation of radiation protection, dose constraints in accordance with Section 9 of the Radiation Act shall be used.* Under section 7 c(5) of the Nuclear Energy Act, *the Radiation and Nuclear Safety Authority shall, to the extent necessary, monitor and oversee the environment of a nuclear facility to verify the reliability of measurements of radioactive releases and to ascertain the environmental impact of the facility.* Environmental radiation monitoring ensures for its own part that the annual dose of an individual in the population, arising from the normal operation of a nuclear power plant, stays below the annual dose constraint of 0.1 millisievert as regulated in Section 22 b of the Nuclear Decree (161/1988). The annual dose constraint is less than 2% of the average annual dose of Finnish people of 5.9 mSv (Siiskonen, 2020).

Radiation exposure arising from the operation of a nuclear power plant shall be kept as low as reasonably achievable. A nuclear facility and its operation shall also be designed so that the constraints presented in the Nuclear Energy Decree are not exceeded. It is not sufficient to stay within the constraints, the releases of radioactive substances and environmental radiation levels resulting from the operation of a nuclear facility shall be kept as low as possible. The holder of a licence entitling to the use of nuclear energy shall derive the release limits of radioactive substances for the nuclear power plant in such a way that the constraint on the individual dose under the Nuclear Energy Decree is not exceeded.

STUK's Guide YVL C.7 gives the detailed requirements applicable to the licensee for the radiological monitoring of the environment of a nuclear facility. The licensee shall draw up a programme for the radiation monitoring of the environment of the nuclear facility and report the results of the programme to STUK. According to Guide YVL C.7, STUK will perform independent regulatory control in the environment of the nuclear facility during the operation of the nuclear facility by collecting and analysing samples from the surroundings of the nuclear facility to a necessary extent. With regard to arranging the environmental monitoring of nuclear facilities, the IAEA has also issued a guide Environmental and Source Monitoring for Purposes of Radiation Protection, IAEA Safety Standards Series No. RS-G-1.8 (IAEA, 2005). The entity formed by the environmental monitoring conducted by the licensee and STUK is in line with the recommendations regarding the contents of the IAEA's monitoring programme.

The results of environmental monitoring conducted by STUK are compiled to this report. The results are compared against the environmental monitoring findings and releases reported by the licensees.

3 Releases from nuclear power plants

During the normal operation of nuclear power plants, radioactive substances are generated, a very small proportion of which may end up in the environment. Radioactive substances are mostly generated to the reactor's nuclear fuel as a result of nuclear fission and as activation products of various materials as a result of neutron radiation. Radioactive substances remain mostly inside the fuel rods as the rod cladding prevents the release of the substances into the surrounding cooling water. The reactor cooling system and related cleaning and waste systems also contain radioactive substances. Gaseous radioactive substances are also generated to the fuel which can, by diffusion, leave the fuel rods. In rare cases, the fuel rod cladding can get damaged in use and lose its tightness, increasing the radioactivity of cooling water.

During the normal operating conditions, the nuclear facility releases into the atmosphere the facility's ventilation exhaust air and the gaseous substances removed from the processes, which have been purified, if necessary. Gaseous releases are directed to the power plants' exhaust air system. Liquid radioactive substances generated at the nuclear power plant are reduced by evaporation, filtration and delaying before they are discharged into the sea. Releases of liquid radioactive substances are discharged with the power plant's cooling water into the sea. In disturbance and accident situations, radioactive substances can be released into the environment also via abnormal routes and the composition of the releases may differ from the releases of normal operation.

In December 2021, the surface water of the Cs separated evaporation concentrate tanks was discharged at the liquid waste storage of the Loviisa power plant. This discharge is carried out approximately once every four years. As a result of the discharge, radioactive releases into the sea were higher than they have been in the years when the discharge has not been made. Iodine releases were also higher than usual as a result of a minor fuel leak at LO2. However, similarly to previous years, the 2021 releases were considerably below the set release limits.

At Olkiluoto, the OL3 plant unit reached its first criticality on 21 December 2021. This plant unit did not yet generate any releases of radioactive substances to be reported.

In 2021, the radioactive discharges from the power plants were small in relation to the set release limits (Fortum, 2022; TVO, 2022). In Loviisa, the release of noble gases into the atmosphere (Kr-87 equivalent release) was approximately 0.04% and the release of iodine (I-131 equivalent release) was approximately 0.025% of their respective release limits in 2021. The release of tritium (H-3) into the sea was approximately 10% and the release of fission and activation products into the sea was approximately 0.11% of their respective release limits.

At Olkiluoto, the release of noble gases into the atmosphere was 0.002% and the release of iodine (at Olkiluoto, the release limit was set for I-131) was 0.12% of their respective release limits. The release of tritium into the sea was approximately 2.2% and the release of fission and activation products into the sea was approximately 0.05% of their respective release limits.

Typical radionuclides originating from the Loviisa power plant and found in the vicinity of the power plant are H-3, Co-60 and Ag-110m and those of the Olkiluoto power plant are H-3, Mn-54, Co-58 and Co-60. The differences in the observed nuclides are due to the different plant types and differences in the materials used in the reactor circuits, for example.

The nuclides causing the largest calculated dose for an individual in the most highly exposed population group are C-14 for air releases and Co-60 or Cs-137 for water releases. The calculated dose estimate for the part of Loviisa C-14 atmospheric releases is based on the conservative assumption that C-14 is fully released into the environment as carbon dioxide (CO₂). However, studies show that 80 – 90% of C-14 releases are released as hydrocarbons that are not easily absorbed by plants (Fortum 2022). For this reason, the samples collected by the environmental monitoring from the surroundings of the nuclear power plants show very little difference to the C-14 activity concentrations of samples collected from elsewhere in Finland. For this same reason, the calculated dose estimate of an individual in the most highly exposed population group is probably also estimated to be higher than it actually is for the part of C-14.

Annex 1 presents in more detail the most common radionuclides detected in releases from the nuclear power plants and in environmental monitoring. Not all radionuclides detected in environmental monitoring originate from the nuclear power plants. There is also natural radioactivity and artificial radionuclides in the environment, such as H-3, Sr-90 and Cs-137, originating from the nuclear weapons testing of the 1950s and 1960s and, in particular, from the Chernobyl disaster of 1986.

4 Environmental monitoring programme of the licensee

The holder of a licence entitling to the use of nuclear energy shall monitor the concentrations of radioactive substances in the environment of the power plant. Guide YVL C.7, published by STUK, provides the minimum requirements for the licensee's environmental radiation monitoring programme (Annex 2):

- The programme shall include external radiation measurements carried out using environmental dosimeters located in the plant's terrestrial environment and external radiation dose rate measuring stations.

- In the terrestrial environment, the measurements shall be primarily focused on the definition of radioactive substances in the air, atmospheric deposition, household water and garden products. In addition, the monitoring programme shall examine the radioactive substances in the indicator organisms in the terrestrial environment. Indicator organisms refer to organisms and plants that collect or enrich radionuclides particularly well and are therefore suitable for the monitoring of radionuclides in the environment.
- In the water environment, the measurements shall be focused on the definition of radioactive substances diluted and mixed in the water.

The results of the environmental radiation monitoring of the licensee are presented in the licensee's annual report for environmental radiation safety, which the licensee submits to STUK (Fortum, 2022; TVO, 2022). STUK assesses the adequacy of the licensee's own monitoring programme and its results and compares the results of the licensee's monitoring with those of STUK's own monitoring programme. The results of the licensee's programme are covered later in Section 6, Results of environmental monitoring, where the results are compared to the results of STUK's monitoring programme.

On 30 December 2021, Posiva Oy submitted to the Finnish Government an operating licence application for a spent nuclear fuel encapsulation plant and final disposal facility. According to Posiva's plan, the environmental radiation monitoring programme during operation will be carried out as part of TVO's environmental monitoring programme. Monitoring targets covering potential Posiva release routes will be added to TVO's monitoring programme.

5 Environmental monitoring programme and methods of the Radiation and Nuclear Safety Authority

STUK's environmental radiation monitoring programme is designed to take into account the conditions of the plant sites and their surroundings and the operation and use of the plants. In this way, the radiation monitoring of the environment is carried out correctly targeted and dimensioned. In addition to the planned programme, additional sampling may be carried out if necessary, for example in the event of plant transients. Environmental radiation monitoring is targeted at the plant environment and the surrounding population.

Measurements are made on the terrestrial and marine environment samples, in addition to which air samples are collected during the annual outages of the plants. Sampling focuses primarily on food chain-related sample types, such as milk, agricultural products, household water, fish, game and other food. In addition, the radiation monitoring programme includes indicator organisms and materials of the aquatic and terrestrial environment, such as wild terrestrial and marine environment flora and sinking matter.

The same or similar sample types are collected from the environment of both nuclear power plants, taking into account local conditions. The sampling items and types are selected so that

they reflect as well as possible the state of the immediate surroundings of the plants. Samples are taken representatively up to a distance of several kilometres from the plant, taking into account any release routes of radionuclides, the dispersion of releases into the environment, the habits of the population and the location of settlement in the environment. The radionuclide concentrations of the samples are compared against the radionuclide concentrations of samples collected elsewhere in Finland and with the observations of previous years.

5.1 Monitored pathways and sampling

The sample types are divided into three main groups: air and terrestrial and marine environment samples. In addition to these, the accumulation of radioactive substances in the inhabitants in the vicinity of the power plant is studied. Sampling of the environmental samples according to the monitoring programme is usually carried out by STUK's sampler. Some samples under the monitoring programme are obtained directly from local farmers, breeders or other operators. The sampling schedule is presented in Annex 3.

In 2021, quality assurance sampling was also included in the programme. Parallel samples will be collected annually from one marine environment and one terrestrial environment target. In 2021, the sample species were seawater and grazing grass.

5.1.1 Outdoor air and atmospheric deposition

Continuous collection of outdoor air samples is part of the licensee's monitoring programme. STUK collects outdoor air particle samples supplementing the licensee's measurements during the annual outages of the plants. STUK's supplementary air sample collector (Figure 1) is equipped with fibreglass filters. The air samplers of the licensees also use active carbon cartridges. The fibreglass filter collects aerosols, which are solid or liquid particles floating in the air. Typical aerosol particles are of a micrometre size. The activated carbon cartridge collects gaseous substances, such as radioactive iodine. The air sampler flow meters measure the air volumes passing through the fibreglass filter and the carbon cartridge. The accumulated radioactivity in the filter and carbon cartridge is calculated in Bq/m³ in proportion to the volume of air pumped through the filter.

The licensees' programme determines gamma-active radionuclides from the atmospheric deposition samples. In addition to these, STUK examines the activity concentration of Sr-90 from the whole year's composite atmospheric deposition samples collected by the licensees as part of STUK's monitoring programme.



Figure 1. Mobile air sample collector. Photo: STUK.

5.1.2 Terrestrial environment

The terrestrial environment samples include soil, reindeer lichen, haircap moss, needles, ferns, mushrooms, berries, game, milk samples, grazing grass, crops, root vegetables, domestic water, groundwater and sludge.

Monitoring of soil radioactivity is carried out as a survey every two years. Samples are collected from the surface layer of soil. Radionuclides can get carried via atmospheric releases of the power plants to surface soil. Soil samples were collected in 2020, but the results are reported in this report (Annex 5). The samples will be collected for the next time in 2022. Surface soil radionuclides can increase the exposure of humans to radioactivity directly by increasing the external radiation dose or indirectly through food. Terrestrial samples are taken from 3 – 5 locations in the surroundings of both power plants every two years. The samples are taken at a depth of 0 – 5 cm, for example with a golf hole cutter (Figure 2) and 3 – 5 primary samples are taken from the same depth to be combined into a single sample. Where appropriate, samples may also be taken to examine the depth profile of radionuclides. A flat, open and intact area with as few stones and roots as possible is selected as the sampling area.

Species of wild plants identified as good enrichers of radioactive substances, such as haircap moss, reindeer lichen, ferns (common polypody (*Polypodium vulgare*) and ostrich fern (*Matteuccia struthiopteris*)) and spruce needles (new annual growth), are collected once a year. Haircap moss could not be collected from Loviisa anymore in 2021, and it was replaced by fern. The plants are dried and ground. Reindeer lichen is picked from an area where there are as few other species and organic debris as possible. In addition to the neighbouring areas of the power plants, reference samples of each species are collected from elsewhere in Finland. Samples of berries and four different species of mushroom are also collected annually from the vicinity of the power plants according to local availability. Efforts are being made to obtain samples of game meat from local hunters for examination.

By examining agricultural products and domestic water, it is possible to assess the exposure of humans to radioactive substances via food. Milk samples from nearby dairies are collected by the dairies in containers supplied and labelled by STUK. The milk comes from dairy farms in the vicinity of the power plants, the longest distances from the power plant being about 40 km. Sampling observes the general procedures for the food sampling of milk. Grazing grass samples are collected from the vicinity of the power plant once during the growing season. The aim has been to select pastures so that the milk from the cows grazing on them goes to the same dairies from where the milk samples of the monitoring programme are collected. The analysis of garden and agricultural products includes different cereals and root vegetables. Crop samples are obtained from local farmers at grain farms located in the environment of the power plants extending to a distance of approximately 20 km. Samples are taken from two varieties of cereal once a year after harvest. One root vegetable sample (potato, carrot, swede) is obtained from local producers in the vicinity of the power plants once during the summer season.

Household water samples are collected from the household water of the cities of Loviisa and Rauma twice a year, in the spring and autumn. As part of STUK's monitoring programme, Sr-90 is also determined from household water samples taken at the power plant, which are covered by the licensees' programme. The groundwater sample is collected from a well producing groundwater near the power plants or directly from the groundwater pipeline. Sludge samples are collected before or after and once during the annual outage from the water treatment plants of the neighbouring cities.



Figure 2. Slicing of a soil sample taken with a golf hole cutter. Photo: STUK.

5.1.3 Marine environment

Marine samples help to monitor the dispersion of power plant releases in the marine environment and their accumulation in marine environment flora and fauna. Seawater, periphyton, bladder wrack, aquatic plants with submerged leaves (spiked water milfoil), bottom

fauna, fish, bottom sediment and sinking matter are collected from the marine environment. The sampling points for the marine environment are shown in Figures 3 and 4.




Figure 3. Sampling points in the marine environment of Loviisa. The map includes data from the Background map series 04/2020 of the National Land Survey of Finland.




Figure 4. Sampling points in the marine environment of Olkiluoto. The map includes data from the Background map series 03/2021 of the National Land Survey of Finland.

Seawater samples (surface water) are collected from several sampling points in the environment of the power plants. The point closest to the nuclear facility is sampled more frequently and the others less frequently.

The examination of aquatic plants includes bladder wrack (*Fucus vesiculosus*) and spiked water milfoil (*Myriophyllum spicatum*). These plants efficiently collect radionuclides from water and are therefore good release indicators. Bladder wrack samples are taken at several points several times a year (Figure 5). Plants are collected both in the areas where the cooling water is discharged and further away from the power plant. Especially in Loviisa, general changes in environmental conditions (e.g. eutrophication of waters) are reflected in a deterioration of the bladder wrack population near the power plant. In spring 2021, a sample meeting the sampling criteria could no longer be collected from the point closest to the Olkiluoto power plant, so the bladder wrack sample was replaced with a sample of spiked water milfoil. You can still find spiked water milfoil on the discharge side of the plant. Also, periphyton is collected as algal samples. Periphyton refers to organisms attached to a solid surface in water, mainly algae.

Periphyton is collected onto a 50 x 50 cm polycarbonate plate throughout the growing season (May–November). In Loviisa, periphyton has also been collected throughout the winter season (November–April). Factors affecting the growth of periphyton include flow rate, light and water quality.



Figure 5. Sampling of bladder wrack by means of scuba diving, with the Olkiluoto power plant in the background (above). Bladder wrack (below). Photos: STUK.

The bottom fauna sample type includes blue mussel or *Saduria entomon*, depending on availability. Blue mussel could not be found from the Iso Kaalonperä sampling point in Olkiluoto, so dark false mussel was collected as a replacement sample. The samples are collected from one sampling point once a year (Figure 6). The fishing of fish samples is done once a year in May–October and the number of sample species must be at least four every year: for example, Baltic herring, pike, perch and bream. As regards Loviisa, a fry sample received from a nearby fish farm is also examined. There is no fry farming activity near Olkiluoto. If necessary, a sample of Baltic herring suitable for analysis may be obtained from a local fish wholesaler, provided that the normal sampling fishing does not yield catch. In addition, a comparison sample of pike is taken, whose fishing area is not in the immediate vicinity of the power plants.

Sinking matter refers to particles that sink in water towards the bottom, consisting mainly of organic solids produced in the open sea area and in the shore zone, organic and inorganic solids brought by runoff water and river water and solid matter of bottom sediment getting occasionally mixed with water. Sinking matter is collected year-round from several sampling points into cylindrical collecting tubes, which are anchored to the desired depth (Figure 7). The bottom sediment samples are collected annually from 5 – 6 points and a surface layer of 0 – 5

cm is taken as a sample. The bottom sediment sample is taken with a dedicated cylindrical sediment collector with a steel structure (Gemini), which sinks into the sediment due to its own weight or additional weights and the closure mechanism locks the sediment plug inside the collector.



Figure 6. A bottom set pot for catching *Saduria entomon* (on the left) and *Saduria entomon* (on the right). Photos: STUK.



Figure 7. A collector of sinking matter. Photo: STUK.

5.1.4 Inhabitants of the surroundings

Once a year, people living in the environment of the nuclear power plant are given the opportunity to participate in a measurement to determine the amount of radioactive substances accumulated in the human body. The aim is to get a minimum of 20 inhabitants in the vicinity of both power plants to take part in the measurement every year by sending them an invitation letter by post. The invitation is sent primarily to persons with a residential address within 7 km of the nuclear power plant in the year of arranging the measurement. The group of invited persons is supplemented by a sample of persons whose residential address is within 10 km of the nuclear power plant. The name and address data are based on the data in the Population

Information Register of the Finnish Digital and Population Data Services Agency. Persons of age are invited to take part in the measurement. Participation in the measurement is voluntary and the measurement results are used in a form that does not allow the results to be associated with individuals or their residential addresses.

The gamma-radiating radionuclides contained in the body of the inhabitants of the surroundings of the nuclear power plant are determined by direct gamma-spectrometric measurement from outside the body. This so-called whole-body counting is carried out with special measuring equipment built on a truck. The measurement takes approximately 15 minutes and, during the measurement, the person sits on a chair inside a background radiation shield. During the measurement, the body is not subjected to radiation and no samples are taken from the subject. The measurements would show if the population in the vicinity would have accumulated deviating quantities of radionuclides originating from the power plant. The person receives their measurement result immediately after the measurement.

5.2 Monitoring methods

Sampling and laboratory analyses under the monitoring programme are primarily carried out by the Environmental Radiation Surveillance and Emergency Preparedness department of STUK. Milk, crops, root vegetable, fry, game and sludge samples come from external suppliers. The C-14 analyses of the monitoring programme are carried out at the Laboratory of Chronology of the University of Helsinki. The Environmental Radiation Surveillance department of STUK is a testing laboratory T167 accredited by FINAS, accreditation requirement EN ISO/IEC 17025:2017.

5.2.1 Sample processing and analysis

The samples are sent to STUK's laboratory in Helsinki for analysis. If necessary, the samples are cleaned to only contain the studied sample type. Spoiled samples and samples otherwise not meeting the quality criteria are rejected in the pre-processing phase.

Food samples are processed so that the measurements are made from the edible parts. A preservative is added to the milk samples to prevent contamination. The milk samples are evaporated under the heat lamps and burnt. The iodine content (I-131) of milk is determined from a separate sample by ion exchange. The samples to be dried (lichen, moss, needles, ferns, mushrooms, grazing grass (berries, game, crops, root vegetables, sludge, fish, bottom fauna, bladder wrack, periphyton, aquatic plants) are dried in a drying oven and, then, homogenised by grinding. Fresh and whole fry are measured. The sediment samples and the sinking matter are dried in a freeze dryer and homogenised by grinding. The soil samples are dried in a drying oven and sifted with a 2-mm sieve. Seawater samples are evaporated to a smaller volume by means of heat lamps. For the radiochemical analyses of strontium, the samples are burnt after a gamma-spectrometric measurement. For the determination of tritium, the water samples are distilled. The results of food and environmental samples are reported per sample volume or dry weight (DW), except for the results of mushrooms, berries, root vegetables, game and fish, which are reported per fresh weight (FW). The activity concentrations of dried samples per unit of weight are significantly higher than those of fresh samples.

Radionuclides emitting gamma radiation are analysed from all samples, including Co-60, I-131, Cs-134 and Cs-137. Radionuclides emitting gamma radiation are identified by the energies of gamma radiation typical of each isotope.

Radiochemical methods are used to analyse the alpha- and beta-active substances (H-3, Sr-90 and Pu-238, Pu-239 and Pu-240) of the samples. The analyses of strontium's short-lived isotope Sr-89 was discontinued in 2021. Sr-89 can still be determined from samples, if another instance of environmental monitoring indicates an abnormal environmental emission. Data from previous years shows that the Sr-89 activity concentrations in environmental samples have remained below the specified limits.

In the radiochemical analysis, the chemical separation of the elemental to be examined is carried out first from the sample. In the determinations of strontium, stable Sr and Cs carrier is first added to the samples and the dry samples are liquefied. Strontium is separated from the sample by an extraction chromatography method and Sr-90 is measured from the sample with a liquid scintillation spectrometer. Strontium yield determination is done by using an inductively coupled plasma-mass spectrometer (ICP-MS). H-3 is determined directly from the distilled water sample by means of a liquid scintillation spectrometer. For plutonium analyses, the Pu-242 tracer is added to the samples and the samples are liquefied before chemical isolation. Plutonium is separated from the other alpha-active radionuclides by ion exchange, and the measurement sample is prepared by precipitation and measured by alpha spectrometry. The resolution of an alpha spectrometer is not sufficient to distinguish plutonium isotopes Pu-239 and Pu-240 from each other and, therefore, the results indicate their combined activity concentration in the samples. C-14 is determined from dried samples at the Laboratory of Chronology of the University of Helsinki.

The radiochemical methods are cumbersome and time-consuming compared to the simple determination of nuclides emitting gamma radiation and, therefore, it is not possible to routinely determine the alpha- and beta-active radionuclides from each sample. Radiochemical analyses have been selected for sample types where they play a significant role in human radiation exposure (e.g. Sr-90 in milk and H-3 in household water) or where they may occur (e.g. H-3 in seawater and Pu-239 or Pu-240 in marine environment sediments). If the results of the monitoring programme were to indicate an increase in the activity concentrations of some alpha- or beta-active radionuclides in the samples, it is possible to increase the scope and frequency of the radiochemical analyses. The analyses to be carried out on the different sample types are shown in Table 1.

Measurement times of the samples vary according to the sample and may, in individual cases, be longer than usual, for example, when a sample is left for measurement for the weekend. A longer measurement time may be the reason for the lower than average detection limit reported for some individual samples. In this case, smaller quantities of radioactive substances may also be detected in individual samples. The calculated activity concentrations of the radionuclides correspond to the average of the collection period and, therefore, the activity concentrations do not accurately reflect the activity concentrations of short-term releases that are temporarily higher. The uncertainty of the results is given with an accuracy of 2σ (95% confidence interval).

Table 1. Monitored pathways and analysed radionuclides in STUK's monitoring program.

Monitoring item	Gamma	Sr-90	C-14	H-3	Pu-238, Pu-239, Pu-240
Outdoor air	x				
Atmospheric deposition (annual sample)		x			
Soil	x	x			
Reindeer lichen	x				
Haircap moss	x				
Needles	x		x		
Ferns	x	x			
Mushrooms	x				
Berries	x				
Game	x				
Milk	x	x			
Grazing grass	x		x		
Crops	x	x			
Root vegetable	x				
Household water	x	x		x	
Groundwater	x				
Sludge	x				
Seawater	x	x		x	
Fry	x				
Periphyton	x				
Bladder wrack	x	x			x
Spiked water milfoil	x				
Bottom fauna	x	x			
Fish	x	x			
Surface sediment	x	x ¹			x ¹
Sinking matter	x				x
Inhabitants of the surroundings	x				

¹Analysis was exceptionally not carried out in 2021

6 Results of environmental monitoring

Approximately 460 samples were collected and analysed in the terrestrial and marine environment of the Olkiluoto power plant during 2021. Approximately 140 of the samples were STUK's monitoring samples and the rest were part of the licensee's own monitoring programme. Approximately 450 samples of the terrestrial and marine environment of the Loviisa power plant were examined during 2021. Of these, 125 were STUK's regulatory oversight samples. In addition to these, the radioactivity accumulated in the bodies of the inhabitants in the surrounding area of both power plants was measured.

The detailed analysis results of STUK's control measurements of the 2021 samples are given in Tables 2 – 22 and in Annex 5. The results of the licensees' own measurements are discussed in the text. All radionuclides present in the result tables do not originate from the Olkiluoto or Loviisa power plants. There is always radioactive isotope of potassium K-40 in the environmental samples and in human beings, forming usually the majority of the natural radioactivity of the samples. The terrestrial environment samples also contain Be-7, which is produced in the upper atmosphere due to cosmic radiation. Almost all samples contain a small amount of radionuclide Cs-137, originating from the nuclear weapon tests conducted in the atmosphere and from the Chernobyl disaster. In addition to K-40, this old Cs-137 forms a part of the background concentration observed in the environmental samples. Typical background concentration of Cs-137 in external air in Finland is around 1 – 4 $\mu\text{Bq}/\text{m}^3$ and the Cs-137 atmospheric deposition in the Helsinki area is typically less than 0.5 Bq/m^2 per month (Mattila and Inkinen, 2021). In the Gulf of Bothnia and the Gulf of Finland, the concentration of Cs-137 in seawater is usually around 20 – 30 Bq/m^3 and the concentration of Sr-90 around 4 – 11 Bq/m^3 (HELCOM, 2018). The Cs-137 background concentration in the terrestrial and marine environment can vary strongly according to geographical location, as has been observed, for example, in the activity concentrations of Cs-137 in the Baltic Sea sediments (HELCOM, 2018). If the monitoring samples were to show Cs-137 originating from the power plant, this could be observed in elevated concentrations compared to the regional background concentration and observations from previous years and in the appearance of another radionuclide Cs-134, with a shorter life, in the environmental samples.

6.1 Outdoor air and atmospheric deposition

Continuous monitoring of radioactive substances in outdoor air is the responsibility of the licensee. STUK carries out outdoor air sampling supplementing the licensee's measurements in conjunction with the annual outages at the plant sites (Table 2). The Cs-137 activity concentration of air samples was low. No radionuclides originating from the power plants were detected in the licensee's own outdoor air samples.

Table 2. An air sample supplementing the licensees' monitoring.

Collection site	Collection period	Co-60 $\mu\text{Bq}/\text{m}^3$	Cs-137 $\mu\text{Bq}/\text{m}^3$	Uncertainty 2σ
Loviisa	30/08 – 03/09/2021	< 0.78	0.75	42%
Olkiluoto	05/05 – 07/05/2021	< 0.69	0.80	38%

The collection and monitoring of atmospheric deposition samples belong also to the monitoring programme of the licensees. The licensee's programme determines gamma-active radionuclides from the atmospheric deposition samples, and Sr-90 is determined from the annual sample combined from these atmospheric deposition samples as part of STUK's monitoring programme. In Loviisa, the total surface activity of Cs-137 of the atmospheric deposition samples of the whole year varied between 0.85 and 2.0 Bq/m², while the tritium concentration remained below 2 Bq/l with the exception of one sample that had an H-3 activity concentration of 2.1 Bq/l. At Olkiluoto, the corresponding range was 0.71 – 1.2 Bq/m², while the tritium concentration remained below 2 Bq/l. No other nuclides originating from the power plants were detected. The activity concentrations of Cs-137 and H-3 observed correspond to those found in samples collected elsewhere in Finland. The Sr-90 results of the composite atmospheric deposition samples for the whole year varied between 0.08 and 0.09 Bq/m² (Table 3), which is at the same level or lower than the Sr-90 concentrations observed in the atmospheric deposition samples of the national environmental radiation monitoring in different cities in Finland (Mattila and Inkinen, 2021).

Table 3. Sr-90 results of the composite annual sample of atmospheric deposition.

Site	Collection period	Sr-90 Bq/m ²	2σ
Loviisa			
LPO	31/12/2020–31/12/2021	0.089	10
Smoltti	31/12/2020–31/12/2021	0.076	10
Olkiluoto			
Weather mast	23/12/2020–22/12/2021	0.083	11

6.2 Terrestrial environment

No nuclides originating from the power plants were detected in terrestrial environment samples (Table 4). The Cs-137 concentrations in the terrestrial environment samples varied between different samples. Moss, fern and lichen effectively collect radionuclides from their environment and in some places, these plants can show even high concentrations of Cs-137 originating mainly from the Chernobyl disaster. In previous years, high Cs-137 activity concentrations have been observed in plants in the terrestrial environment of Loviisa, and this year's results do not differ from those of previous years. The activity concentrations of Cs-137 detected in ferns (Figure 8) originate at least mainly from the fallout caused by the Chernobyl nuclear power plant accident in 1986. The emission originating from the nuclear power plant also includes shorter-lived Cs-134 with the half-life of 2.06 years. Small amounts of Cs-134

were detected in fern samples up until the early 2010s. Based on the activity concentrations of the isotopes, they probably originated from Chernobyl. Even though higher concentrations have since been observed again, Cs-134 has not been detected anymore. Based on this, the higher activity concentrations are probably not from a new and more recent source. The ferns' activity concentrations in relation to Cs-137 also varied greatly even within very small distances. Good examples of this are the samples collected in 2021 in Loviisa within a couple of kilometres from each other with an approximately 10-time difference in their activity concentrations. At Olkiluoto, the sampling place has been varied as a result of poor growth of common polypody and in 2018, the sampling species had to be changed to ostrich fern.




Figure 8. Cs-137 activity concentrations [Bq/kg per dry weight] of fern samples collected from Loviisa and Olkiluoto in 2005 – 2021.

The licensee collected fern samples from the Olkiluoto surroundings. Only natural nuclides and Cs-137 were detected in this sample, and the concentration of Cs-137 was 85 Bq/kg, which was low compared to the activity concentrations measured in STUK's monitoring programme samples. In the surroundings of the Loviisa power plant, the licensee collected a sample of ferns (*Dryopteris carthusiana*), which showed no nuclides originating from the power plant.

Table 4. Monitoring measurement results of the lichen, moss, needle and fern samples in 2021. Haircap moss could not be collected from Loviisa and it was replaced by fern.

Site	Collection date	Be-7 Bq/kg	2σ	Cs-137 Bq/kg	2σ	C-14 Bq/kg	2σ	Sr-90 Bq/kg	2σ
Spruce needles									
Loviisa	26/05/2021	17	18%	420	10%	121	5%	-	
Olkiluoto	03/06/2021	7.2	25%	170	17%	128	5%	-	
Reference Kimola	07/06/2021	4.6	23%	61	9.2%	121	5%	-	

6 RESULTS OF ENVIRONMENTAL MONITORING

Reindeer lichen									
Loviisa	03/08/2021	260	9.3%	223	9.1%	-	-	-	-
Olkiluoto	03/06/2021	176	8.0%	151	7.7%	-	-	-	-
Reference Kimola	07/06/2021	243	12%	189	10%	-	-	-	-
Haircap moss									
Loviisa	replaced by fern								
Olkiluoto	27/07/2021	240	10%	378	10%	-	-	-	-
Reference Kimola	10/10/2021	471	10%	775	10%	-	-	-	-
Fern									
Loviisa Hästholmen (common polypody)	18/11/2021	160	11%	3,300	10%	-	-	-	-
Loviisa Källa (Common polypody)	31/08/2021	140	8.2%	330	7.6%	-	6.5	8.9%	
Reference, litti (common polypody)	08/10/2021	105	8.6%	590	4.6%	-	15	8.8%	
Olkiluoto (ostrich fern)	27/07/2021	140	10%	110	10%	-	4.1	9.3%	
Reference Kimola (ostrich fern)	04/10/2021	260	10%	1,200	9.1%	-	17	8.8%	

The Cs-137 activity concentrations of mushroom samples collected in the vicinity of the Loviisa and Olkiluoto power plants varied between 17 and 670 Bq/kg per fresh weight (Table 5). Following the Chernobyl fallout, mushroom samples may occasionally show elevated concentrations of Cs-137 and, in the case of specific mushroom species (such as milk caps), it is common to find exceedances of 600 Bq/kg also in areas of minor fallout (Kostiainen and Ylipieti, 2010). The EU-recommended limit value for natural food placed on the market is 600 Bq/kg (Commission Recommendation 2003/274/EC). Berry samples collected in the vicinity of the power plants and game meat samples (elk meat) obtained from local hunters had low activity concentrations of less than 50 Bq/kg per fresh weight. No game samples could be collected in the Olkiluoto area in 2021.

Table 5. Monitoring measurement results of the mushroom, berry and game samples in 2021. No game samples could be collected in Olkiluoto in 2021.

Site	Collection date	Species	Cs-137 Bq/kg FW	2 σ
Mushrooms				
Loviisa	19/08/2021	rufous milk cap	108	11%
	01/09/2021	velvet bolete	235	8.0%
	01/09/2021	Cortinarius caperatus	379	16%
	07/09/2021	woolly milk cap	355	12%
	07/09/2021	cep	124	16%
Olkiluoto	26/08/2021	cep	40	10%
	26/08/2021	chanterelle	17	10%
	26/08/2021	pickle milk cap	169	7.1%
	14/09/2021	Cortinarius caperatus	175	8.0%
	16/09/2021	rufous milk cap	671	16%
Berries				
Loviisa	06/09/2021	lingonberry	9.6	17%
Olkiluoto	14/09/2021	lingonberry	2.7	23%
	11/08/2021	Sea buckthorn	< 0.04	
Game				
Loviisa	15/11/2021	elk	48	16%
Olkiluoto	-	-		

The Cs-137 activity concentrations of milk samples varied between 0.05 and 0.50 Bq/l (Table 6) in the environmental radiation monitoring programmes of the nuclear power plants. The Sr-90 activity concentration of the milk sample combined from samples for the whole year (20 – 40 km from the power plants) was 0.02 Bq/l in the Loviisa sample and 0.03 Bq/l in the Olkiluoto sample. The activity concentrations are well in line with the activity concentrations of the national environmental monitoring of milk samples, which were between 0.11 and 1.4 Bq/l for Cs-137 and between 0.02 and 0.03 Bq/l for Sr-90 in 2020 (Mattila and Inkinen, 2021). Figures 9 and 10 show the Cs-137 activity concentration in milk samples of the environmental monitoring programmes of the nuclear power plants in 2010 – 2021. Every other month, I-131 was also screened from the samples delivered from dairy farms within a distance of 20 km from the power plants. I-131 was not detected in any of the milk samples (limit of determination 0.007 – 0.018 Bq/l).

In the monitoring programme, the monitoring measurements of the terrestrial environment agricultural products (crops and root vegetable) and grazing grass showed no radionuclides originating from the power plants (Table 7). The Cs-137 activity concentration of the samples was low. The Sr-90 activity concentration of crop samples was 0.12 Bq/kg in Loviisa and 0.26 Bq/kg in Olkiluoto. The C-14 content of grazing grass was 75 – 88 Bq/kg in Loviisa,

63 – 110 Bq/kg in Olkiluoto and 119 Bq/kg in the reference sample (Jokioinen). No radionuclides originating from the power plants were detected in the apple sample from the surroundings of Loviisa in the monitoring programmes of the licensees. The lettuce sample could not be collected from Olkiluoto due to drought.




Figure 9. The Cs-137 concentration (Bq/l) of the milk samples supplied by dairies in the surroundings of the Loviisa power plant (distance of the farms from the plant 0 – 20 km or 20 – 40 km) in 2010 – 2021.




Figure 10. The Cs-137 concentration (Bq/l) of the milk samples supplied by dairies in the surroundings of the Olkiluoto power plant (distance of the farms from the plant 0 – 20 km or 20 – 40 km) in 2010 – 2021.

Table 6. Results of the radioactivity monitoring of the milk samples from the dairies in the surroundings of the Loviisa and Olkiluoto nuclear power plants in 2021.

Locality	Collection period	0–20 km				20–40 km			
		K-40 Bq/l	2 σ	Cs-137 Bq/l	2 σ	K-40 Bq/l	2 σ	Cs-137 Bq/l	2 σ
Loviisa	03/01/2021– 24/01/2021	47	15%	0.05	23%	49	15%	0.05	22%
	07/02/2021– 28/02/2021	46	27%	0.08	15%	48	12%	0.06	17%
	07/03/2021– 28/03/2021	47	15%	0.06	19%	49	15%	0.05	22%
	04/04/2021– 25/04/2021	52	15%	0.08	16%	50	12%	0.05	16%
	02/05/2021– 30/05/2021	49	15%	0.06	14%	48	17%	0.11	18%
	06/06/2021– 27/06/2021	50	27%	0.08	16%	54	23%	0.21	23%
	04/07/2021– 01/08/2021	49	11%	0.07	15%	46	15%	0.11	14%
	02/08/2021– 29/08/2021	48	15%	0.08	16%	47	15%	0.18	17%
	05/09/2021– 26/09/2021	49	15%	0.07	20%	51	13%	0.29	10%
	03/10/2021– 31/10/2021	47	16%	0.06	25%	47	15%	0.08	17%
	07/11/2021– 28/11/2021	47	18%	0.09	20%	49	11%	0.09	15%
	29/11/2021– 26/12/2021	48	10%	0.11	12%	46	17%	0.11	18%
Olkiluoto	03/01/2021– 31/01/2021	49	16%	0.22	21%	49	15%	0.43	11%
	07/02/2021– 28/02/2021	47	11%	0.22	11%	48	12%	0.40	14%
	07/03/2021– 28/03/2021	50	14%	0.16	15%	49	13%	0.45%	10%
	04/04/2021– 25/04/2021	47	15%	0.11	18%	56	19%	0.50	11%
	02/05/2021– 30/05/2021	51	15%	0.14	16%	48	15%	0.37	16%
	06/06/2021– 27/06/2021	54	23%	0.17	24%	53	23%	0.44	23%
	04/07/2021– 01/08/2021	50	10%	0.14	9.0%	49	15%	0.35	10%
	02/08/2021– 29/08/2021	49	16%	0.16	22%	45	9.2%	0.23	8.5%
	05/09/2021– 26/09/2021	49	9.3%	0.08	14%	50	11%	0.34	7.3%

	03/10/2021– 31/10/2021	49	16%	0.10	23%	50	15%	0.37	16%
	07/11/2021– 28/11/2021	50	15%	0.22	17%	47	15%	0.31	12%
	29/11/2021– 26/12/2021	55	15%	0.19	16%	52	15%	0.33	12%

Table 7. Monitoring measurement results of the grazing grass, crops and root vegetable samples in 2021. Root vegetables samples could not be collected from Loviisa in 2021.

Site	Collection date	Species	K-40 Bq/kg	uncertainty 2 σ	Cs-137 Bq/kg	uncertainty 2 σ
Grazing grass						
Loviisa	10/06/2021	grazing grass	1010	15%	0.28	26%
	10/06/2021*	grazing grass	1010	15%	0.35	51%
Olkiluoto	15/06/2021	grazing grass	870	15%	1.6	20%
	15/06/2021*	grazing grass	740	17%	0.88	30%
Reference, Jokioinen	30/06/2021	grazing grass	595	16%	<0.21	
Crops						
Loviisa	15/11/2021	wheat	140	13%	0.30	18%
	15/11/2021	barley	160	12%	0.30	23%
Olkiluoto	06/10/2021	wheat	140	15%	0.29	25%
	06/10/2021	oat	130	13%	0.92	14%
Reference, Kimola	10/10/2021	wheat	130	17%	0.96	17%
	10/10/2021	oat	140	17%	0.92	19%
Root vegetable						
Loviisa	-	-				
Olkiluoto	05/10/2021	potato	130	23%	<0.06	

*Quality assurance sample

The H-3, Sr-90 and Cs-137 activity concentrations in the household water of the cities of Rauma and Loviisa and the Sr-90 activity concentrations in the household waters of the power plants supplied by the licensees were at the same level as household water concentrations elsewhere in Finland (Table 8). The monitoring programmes of the licensees determined the radionuclides emitting gamma radiation in the household water of the power plants four times a year. No radionuclides originating from the power plants were detected in the household water of the power plants. The H-3 activity concentrations of all household water samples were less than 2 Bq/l. The concentrations correspond to the H-3 concentrations measured in household water elsewhere in Finland.

Table 8. Monitoring measurement results of the household water of the cities of Rauma and Loviisa in 2021.

Site	Collection date	H-3 Bq/l	2 σ	Sr-90 Bq/m ³	2 σ	Cs-137 Bq/m ³	2 σ
Loviisa	12/05/2021	< 1.0		-		< 0.19	
	23/09/2021	< 1.0		-		< 1.3	
	12/05/2021– 23/09/2021	-		1.3	11%	-	
Loviisa power plant	02/03/2021– 29/11/2021*	-		2.9	9.2%	-	
Rauma	07/05/2021	< 1.0		-		1.1	23%
	06/10/2021	1.2	61%	-		1.1	25%
	07/05/2021– 06/10/2021	-		4.6	9.2%	-	
Olkiluoto power plant	02/02/2021– 22/09/2021*	-		5.4	9.0%	-	

*Combined annual sample, only Sr-90 to be determined

No artificial radionuclides were found in the groundwater sample taken in the surroundings of Olkiluoto and the Cs-137 activity concentration of the Loviisa groundwater sample was very low, 0.35 Bq/m³. Two sludge samples were taken at the Vårdö wastewater treatment plant near Loviisa, one in February and the other during the annual outage in September. The following artificial radionuclides were detected in the samples: I-131, Lu-177, Tl-201 and Tl-202. In Olkiluoto, one sludge sample was taken in February and another during the annual outage in May at UPM Rauma's wastewater treatment plant. I-131 was detected in the samples. These same radionuclides are also detected in the samples taken annually from the Viikinmäki wastewater treatment plant in Helsinki, and they are common nuclides used in radionuclide therapy in hospitals, for which reason they are likely to have originated from patients who have received radionuclide therapy. In Loviisa, sludge samples are examined in the licensee's own programme at the wastewater treatment plant at the Loviisa power plant site four times a year, during annual outages and outside these times. The following artificial radioactive substances originating from the power plant were found in these sludge samples: Mn-54, Co-58, Co-60, Ag-110m, Sb-124 and I-131. At Olkiluoto, the licensee collected a water sample from the aeration tank of the Olkiluoto landfill, and this sample contained a small amount of Co-60 (3 Bq/m³).

6.3 Marine environment

Some radionuclides originating from the power plants were detected from the samples collected from the marine environment of the Loviisa and Olkiluoto power plants. However, the concentrations of radionuclides were small and insignificant in view of the radiation exposure of the environment.

Tables 9 and 10 show the monitoring measurement results of the seawater samples in 2021. The results are presented in an order of distance from the discharge opening, showing the results closest to the discharge opening at the beginning of the tables.

The seawater samples collected from the surroundings of both power plants showed activity concentrations of H-3 exceeding 2 Bq/l, but the concentrations remained under 10 Bq/l, with the exception of two Olkiluoto samples. The typical concentration of tritium was 1 – 2 Bq/l in seawater of the Baltic Sea area in 2011 – 2015 (HELCOM 2018). Based on the long-term

results of the Baltic Sea area, the background level for the tritium concentration is less than 2 Bq/l for the seawater, rainwater and household water samples in the environmental radiation monitoring of the Olkiluoto and Loviisa power plants. Tritium concentrations above this background level are attributed to releases from the power plants. Small amounts of Co-60 and Mn-54 were also detected in one of the Olkiluoto seawater samples. According to the power plant's report, the OL2 plant unit had discharged water in the morning of the sampling (17/02/2021), and the activity ratios in the seawater sample were almost identical to those of the discharge water. The Sr-90 and Cs-137 activity concentrations in the seawater sample were at a usual Baltic Sea level.

In the longer time series (Figures 11 and 12), it can be seen that the most significant source of Cs-137 in seawater is the Chernobyl disaster in 1986. The effect of normal releases from the plants cannot be distinguished from the activity originating from the Chernobyl disaster as the Cs-137 concentrations of seawater correspond to the common activity concentration of Cs-137 in the Baltic Sea (HELCOM 2018).

The Cs-137 concentrations of the seawater samples taken by the licensees corresponded to the common activity concentration of Cs-137 in the Baltic Sea. The concentration of H-3 in seawater was in the licensee's measurements between 1.79 and 80 Bq/l in Loviisa and between 1.06 and 3.10 Bq/l in Olkiluoto. The high H-3 activity concentration in the seawater sample taken in Loviisa in the spring is explained by the fact that the sample was incorrectly taken at the same time with the process water discharge. In other respects, the analysis results of the seawater samples taken by the licensees corresponded to the results of the samples taken by STUK.

Table 9. Monitoring measurement results of the Loviisa seawater samples in 2021.

Site	Collection date	H-3 Bq/l	uncertainty 2σ	Sr-90 Bq/m ³	uncertainty 2σ	Cs-137 Bq/m ³	uncertainty 2σ
Halkokari 02	12/02/2021	1.6	44%	5.5	9.7%	11	11%
	22/04/2021	2.8	27%	6.0	9.5%	14	14%
	06/08/2021	1.8	38%	5.6	9.6%	13	16%
	18/11/2021	2.8	28%	5.8	9.4%	14	15%
Klobbfjärden 1	21/04/2021	8.5	14%	-		13	12%
	17/11/2021	1.7	41%	-		13	12%
Vådholmsfjärden 4	21/04/2021	1.3	50%	-		12	12%
	16/11/2021	< 1,1		-		15	12%
Hudöfjärden 8	22/04/2021	1.3	50%	-		14	14%
	16/11/2021	< 1,1		-		14	15%
Påsalöfjärden R1	21/04/2021	1.1	60%	4.7	9.7%	9.1	18%
	16/11/2021	< 1,1		5.3	9.6%	11	15%
	16/11/2021*	-		-		11	15%

*Quality assurance sample

Table 10. Monitoring measurement results of the Olkiluoto seawater samples in 2021.

Site	Collection date	H-3 Bq/l	uncertainty 2σ	Sr-90 Bq/m ³	uncertainty 2σ	Cs-137 Bq/m ³	uncertainty 2σ
Iso Kaalonperä 13	17/02/2021*	17	11%	5.8	9.5%	18	10%
	16/04/2021	< 0.97		6.2	9.7%	18	13%
	28/07/2021	1.1	59%	1.5	11%	20	10%
	28/10/2021	32	10%	5.6	9.4%	20	13%
Liponluoto 2	13/04/2021	< 1.0		-		18	14%
	26/10/2021	< 1,1		-		20	11%
Rääpinkivet 3	14/04/2021	< 1.0		-		18	12%
	27/10/2021	< 1,1		-		21	9.8%
Santakari 15	14/04/2021	< 1.0		-		15	17%
	27/10/2021	<1,1		-		18	14%
Kylmäpihlaja 17	15/04/2021	< 0.97				19	14%
	27/10/2021	< 1,1				20	16%
Viikari 16	14/04/2021	< 1.0		6.3	9.8%	16	12%
	27/10/2021	< 1,1		5.6	9.4%	19	16%
	27/10/2021**			-		21	13%

*Co-60 (8.0 Bq/m³) and Mn-54 (1.0 Bq/m³) were also detected in the sample

**Quality assurance sample




Figure 11. The Cs-137 activity concentration in seawater at the nearest (02/A, blue) and furthest (R1, yellow) sampling point of the Loviisa power plant in 1980 – 2021 presented on a logarithmic scale.




Figure 12. The Cs-137 activity concentration in seawater at the nearest (Iso-Kaalonperä 13, blue) and furthest (Santakari 15, yellow) sampling point of the Olkiluoto power plant in 1981 – 2021 presented on a logarithmic scale.

Fishing in the vicinity of the power plants takes place at two different distances from the power plants: in Loviisa the fishing areas are 0 – 2 km and 2 – 10 km from the plant and in Olkiluoto 0 – 3 km and 3 – 10 km from the plant. The Baltic herring is fished at the distance of 0 – 10 km. The Cs-137 concentrations of fish samples (Baltic herring, perch, pike and roach) varied between 1.6 and 10 Bq/kg (per fresh weight, Table 11). The concentrations were low and well in line with the Cs-137 activity concentrations in the fish and reference samples of the Baltic Sea area (HELCOM 2018, Mattila and Inkinen, 2020). The concentrations of Sr-90 in the fish samples were also low. The Cs-137 activity concentration in the fry sample received from the Smoltti fish farm in Loviisa was very low.

The radioactivity concentrations of the bottom fauna samples (*Saduria entomon* and blue mussel) were low (Table 12). The *Saduria entomon* sample from the surroundings of Loviisa contained Ag-110m originating from the power plant, but the concentration was low and does not affect the radiation exposure of the organism.

Table 11. Monitoring measurement results of the fish samples from the marine environment of Loviisa and Olkiluoto in 2021.

Sample type	Site	Collection period	Cs-137 Bq/kg	Uncertainty 2 σ	Sr-90 Bq/kg	Uncertainty 2 σ
Baltic herring	Loviisa 0–10 km	30/04/2021	1.6	11%	0.040	16%
	Olkiluoto 0–10 km	05/05/2021	2.2	9.6%	0.022	21%
Perch	Loviisa 0–2 km	25/05/2021	8.6	13%	0.010	29%
	Loviisa 2–10 km	09/06/2021–06/08/2021	8.1	17%	-	
	Olkiluoto 0–3 km	05/05/2021–19/05/2021	10	16%	0.007	40%
	Olkiluoto 3–10 km	11/08/2021	8.8	8.4%	-	
Pike	Loviisa 0–2 km	08/06/2021–09/06/2021	6.5	18%	-	
	Loviisa 2–10 km	11/06/2021	5.0	12%	-	
	Olkiluoto 0–3 km	05/05/2021	6.8	17%	-	
	Olkiluoto 3–10 km	01/06/2021–28/07/2021	7.2	9.9%	-	
	Pellinki Reference	08/09/2021	6.0	17%	-	
Bream	Loviisa 0–2 km	28/05/2021–10/09/2021	1.7	14%	-	
	Loviisa 2–10 km	26/05/2021–09/09/2021	1.8	10%	-	
	Olkiluoto 0–3 km	05/05/2021–28/10/2021	2.1	14%	-	
	Olkiluoto 3–10 km	20/05/2021–20/05/2021	2.5	23%	-	
Fry sample	Loviisa, Smoltti	09/11/2021	0.49	15%	-	

Table 12. Monitoring measurement results of the bottom fauna collected from the marine environment of Loviisa and Olkiluoto in 2021.

Sample type	Site	Collection period	Sr-90 Bq/kg	2 σ	Ag-110m Bq/kg	2 σ	Cs-137 Bq/kg	2 σ
Saduria entomon	Hästholsfjärden 3	16/05/2021	9.4	6.9%	1.3	25%	6.8	24%
Dark false mussel	Iso Kaalonperä	05/10/2021	7.1	12%	< 0.67		1.2	29%
Blue mussel	Reference, Iso-Pietari C	16/09/2021	9.4	11%	< 0.82		0.62	25%

In the aquatic environment, periphyton, bladder wrack and spiked water milfoil, from among the aquatic plants with submerged leaves, have proven to be particularly good indicators of power plant releases. The longest observation series are of bladder wrack, and they clearly show the impact of power plant releases. However, bladder wrack is sensitive to environmental

changes, and collecting it has not been possible every year. Figures 13 and 14 show the annual averages of the activity concentrations of some of the most significant nuclides originating from the power plant in bladder wrack samples collected nearest the power plant. In the bladder wrack samples, the activity concentrations of the nuclides originating from the power plants have decreased clearly as the power plant releases have decreased. Figures 15 and 16 show the link between the activity concentration of Co-60 in the bladder wrack samples and the Co-60 discharges from the power plant into the sea. Changes in the activity concentrations follow quite closely the changes in the releases, there seems to be a delay of approximately one year in the change in the activity concentrations in the surroundings of the Loviisa power plant. Every four years, the Loviisa power plant carries out releases of the surface water of the Cs separated evaporation concentrate tanks, which cause, for example, an increase in Co-60 releases. These discharges are scheduled for the end of the year in order to mitigate the effects of nutrients contained in the release. Radioactive substances resulting from the release will therefore only be visible in the results of the monitoring of samples for the next growing season. The latest discharge was made in December 2021, and it will most likely be reflected in the 2022 monitoring samples.




Figure 13. Averages of the activity concentrations of the most significant radionuclides originating from the power plant in bladder wrack (Bq/kg per dry weight) at the nearest sampling point of the Loviisa power plant in 1977 – 2021 on a logarithmic scale.




Figure 14. Averages of the activity concentrations of the most significant radionuclides originating from the power plant in bladder wrack (Bq/kg per dry weight) at the nearest sampling point of the Olkiluoto power plant in 1977 – 2021 on a logarithmic scale.




Figure 15. The Co-60 releases into the sea and the average of the Co-60 activity concentration in the bladder wrack samples of the nearest sampling point of the Loviisa power plant in 1977 – 2021.




Figure 16. The Co-60 releases into the sea and the average of the Co-60 activity concentration in the bladder wrack samples of the nearest sampling point of the Olkiluoto power plant in 1977 – 2021.

The results of the periphyton samples are given in Annex 4. Several radionuclides originating from the power plants were found in the periphyton samples, but their concentrations were low. The nuclides originating from the power plants found in the periphyton samples of Loviisa were Mn-54, Co-58, Co-60, Nb-95, Zr-95, Ag-110m, Sb-124 and Cs-137, and those of Olkiluoto were Mn-54, Co-60 and Cs-137. In addition to periphyton samples, Co-60 was also detected in bladder wrack in Olkiluoto (Tables 13 and 14). No radionuclides originating from the power plants were detected in bladder wrack in Loviisa. The Cs-137 activity concentrations of bladder wrack varied between 9.4 – 15 Bq/kg. A second bladder wrack sample could not be obtained from the Iso Kaalonperä sampling point in Olkiluoto due to the lack of plant life; a spiked water milfoil sample was used as a replacement. The same nuclides originating from the power plants have been observed in the periphyton and bladder wrack samples as in the previous years and the activity concentrations do not differ from the those measured in the previous years. The nuclides found in the samples are the same ones that the power plants have reported to have discharged into seawater on the basis of their own release measurements.

Small quantities of radionuclides originating from the power plants were observed in spiked water milfoil (Tables 15 and 16). Co-58, Co-60, Ag-110m, and Sb-124 were detected in the spiked water milfoil sample collected from the discharge area of the Loviisa power plants and Co-60 was detected in the spiked water milfoil sample collected from the Olkiluoto discharge area. The Cs-137 activity concentrations of the spiked water milfoil samples varied between 2.1 and 30 Bq/kg. The reference samples of spiked water milfoil were collected further away from the discharge opening but still in the vicinity of the power plant (Strömslandet in Loviisa and Aikonmaa in Olkiluoto), and radionuclides originating from the power plants were no longer found in these samples, indicating that the activity concentrations of the radionuclides are lower further away from the discharge opening.

Table 13. Radionuclides found in the bladder wrack samples collected in the marine environment of Loviisa in 2021.

Sampling point	Collection date	Cs-137 Bq/kg	2 σ	Sr-90 Bq/kg	2 σ
Stenöarna	11/05/2021	13	11%	5.6	10%
	01/09/2021	10	11%	-	
Hästholmen SW	01/07/2021	11	15%	-	
	01/09/2021	9.4	10%	-	
Lilla Djupberget C	11/05/2021	11	9.4%	-	
	31/08/2021	12	11%	-	
Boistö D	12/05/2021	10	14%	-	
	31/08/2021	13	9.3%	-	
Storskarven E	12/05/2021	10	10%	4.1	10%
	31/08/2021	8.4	15%	-	

Table 14. Radionuclides found in the bladder wrack samples collected in the marine environment of Olkiluoto in 2021. A second sample of bladder wrack could not be collected from the Iso Kaalonperä sampling point.

Site	Collection date	Co-60 Bq/kg	2 σ	Cs-137 Bq/kg	2 σ	Sr-90 Bq/kg	2 σ
Iso Kaalonperä 9	Replaced with spiked water milfoil						
	12/08/2021	0.88	14%	15	14%	-	
Kalliopöytä B	05/05/2021	0.44	19%	11	10%	5.1	
	14/09/2021	0.66	23%	14	11%		
Reimarkrunni	05/05/2021	0.34	17%	9.8	11%	-	
	15/09/2021	< 0.32		9.6	24%	-	
Iso-Siiliö D	04/05/2021	< 0.15		9.0	10%	-	
	16/09/2021	< 0.17		6.7	11%		
Iso-Pietari C	04/05/2021	< 0.15		9.2	14%	-	
	16/09/2021	< 0.19		8.5	15%	-	
Kylmäpihlaja 17	05/05/2021	< 0.11		7.7	14%	-	
	15/09/2021	< 0.14		7.1	11%	-	
Viikari 16	04/05/2021*	< 0.10		11	9.2%	4.5	
	16/09/2021	< 0.22		10	10%	-	

*I-131 (0.21 Bq/kg) was also detected in the sample

Table 15. Radionuclides found in the spiked water milfoil samples collected in the marine environment of Loviisa in 2021.

Site	Collection date	Co-58 Bq/kg	2 σ	Co-60 Bq/kg	2 σ	Ag-110m Bq/kg	2 σ	Sb-124 Bq/kg	2 σ	Cs-137 Bq/kg	2 σ
Halkokari	01/09/2021	0.32	17%	0.42	26%	7.4	15%	0.38	25%	30	10%
Kasabergs- udden	21/09/2021	< 0.47		< 0.62		< 0.56		< 0.47		28	18%
Strömslandet (Reference)	22/09/2021	< 0.32		< 0.39		< 0.37		< 0.34		23	

Table 16. Radionuclides found in the spiked water milfoil samples collected in the marine environment of Olkiluoto in 2021.

Site	Collection date	Co-60 Bq/kg	2 σ	Cs-137 Bq/kg	2 σ
Iso Kaalonperä	29/06/2021	1.2	25%	2.1	20%
	14/09/2021*	0.56	30%	9.3	24%
	14/09/2021	1.4	39%	21	9.9%
Aikonmaa (Reference)	05/10/2021	< 0.60		10	12%

*Replaces a bladder wrack sample

Small concentrations of radionuclides originating from the power plants were found in the sinking matter samples collected from the environment of the power plants (Tables 17 and 18). Co-60, Ag-110m and Cs-137 were detected in Loviisa, and Co-60, Sb-125 and Cs-137 were detected in Olkiluoto. Cs-137 found in the sinking matter originates largely from the Chernobyl disaster. Table 19 presents the activity concentrations of the plutonium isotopes Pu-238, Pu-239 and Pu-240 detected in the bladder wrack samples and sinking matter. Sediment and sinking matter show small background concentrations of the Pu-238, Pu-239 and Pu-240 radionuclides originating from the global fallout from the atmospheric nuclear weapons tests. The activity concentrations of Pu-238, Pu-239 and Pu-240 in the samples from the surroundings of the Olkiluoto and Loviisa power plants are at the same level as those commonly found in sediment in the Baltic Sea region (HELCOM, 2018).

Table 17. Radionuclides observed in the sinking matter samples collected from the marine environment of Loviisa in 2021.

Site	Collection period	Co-60 Bq/kg	2 σ	Ag-110m Bq/kg	2 σ	Cs-137 Bq/kg	2 σ
Hästholsfjärden 5S	12/11/2020– 22/04/2021	0.66	29%	5.1	20%	302	16%
	22/04/2021– 22/06/2021	< 1.2		4.1	22%	173	10%
	22/06/2021– 17/08/2021	< 1.7		4.9	33%	170	13%
	18/08/2021– 18/11/2021	1.3	20%	3.8	18%	259	9.6%
	10/11/2020– 22/04/2021	< 0.54		3.5	23%	281	10%
Hästholsfjärden 3	22/04/2021– 23/06/2021	< 1.4		< 1.6		174	10%

6 RESULTS OF ENVIRONMENTAL MONITORING

	23/06/2021– 30/08/2021	< 1.8		< 3.0		220	11%
	19/08/2021– 17/11/2021	0.52	48%	1.4	31%	235	10%
Klobbfjärden 1	10/11/2020– 22/04/2021	< 1.3		< 1.7		325	8.6%
	21/04/2021– 22/06/2021	< 0.92		1.4	24%	277	16%
	22/06/2021– 18/08/2021	< 1.9		< 2.5		297	12%
	18/08/2021– 17/11/2021	< 0.33		< 0.40		313	6.4%
Vådholmsfjärden 4	10/11/2020– 22/04/2021	< 0.95		< 1.4		290	10%
	21/04/2021– 22/06/2021	< 0.72		< 0.98		129	11%
	23/06/2021– 17/08/2021	< 2.9		< 4.1		196	13%
	17/08/2021– 16/11/2021	< 0.29		< 0.46		281	12%
Påsalöfjärden R1	11/11/2020– 21/04/2021	< 0.83		< 1,1		269	8.6%
	21/04/2021– 23/06/2021	< 0.74		< 1,1		207	11%
	23/06/2021– 17/08/2021	< 1.3		< 1.5		190	10%
	17/08/2021– 16/11/2021	< 0.29		< 0.40		227	16%

Table 18. Radionuclides observed in the sinking matter samples collected from the marine environment of Oikiluoto in 2021.

Site	Collection period	Co-60 Bq/kg	2σ	Sb-125 Bq/kg	2σ	Cs-137 Bq/kg	2σ
Räापinkivet 3	17/11/2020– 14/04/2021	2.4	18%	< 0.92		160	11%
	14/04/2021– 16/06/2021	0.73	42%	< 0.42		113	11%
	16/06/2021– 24/08/2021	< 1.4		< 1.4		132	15%
	24/08/2021– 26/10/2021	1.6	19%	< 0.47		134	12%
Vähä Kivikkokari 12	27/11/2020– 13/04/2021	3.5	14%	< 0.63		158	12%
	13/04/2021– 16/06/2021	1.4	33%	< 0.64		133	11%
	16/06/2021– 24/08/2021	1.2	24%	< 1,1		< 0.86	
	24/08/2021– 26/10/2021	0.78	23%	< 0.36		153	12%

6 RESULTS OF ENVIRONMENTAL MONITORING

Kaalonperä 9	17/11/2020– 15/04/2021	15	7%	1.9	31%	151	7%
	15/04/2021– 16/06/2021	4.3	11%	< 0.48		130	7%
	16/06/2021– 24/08/2021	2.3	28%	< 0.87		144	7%
	25/08/2021– 26/10/2021	1.6	19%	< 0.60		159	13%
Santakari 15	18/11/2020– 14/04/2021	1.0	45%	< 0.64		141	10%
	14/04/2021– 17/06/2021	< 0.49		< 0.41		128	13%
	17/06/2021– 24/08/2021	< 1.7		< 1.8		142	15%
	24/08/2021– 27/10/2021	< 0.47		< 0.43		145	10%
Kuuskajaskari 20	17/11/2020– 15/04/2021	< 1.5		< 1.4		140	15%
	15/04/2021– 17/06/2021	3.4	17%	< 0.52		109	10%
	17/06/2021– 24/08/2021	< 1.8		< 1.5		163	12%
	25/08/2021– 28/10/2021	< 0.47		< 0.45		147	10%
Keskivedenkari 18	18/11/2020– 14/04/2021	< 0.59		< 0.53		136	11%
	14/04/2021– 17/06/2021	< 0.39		< 0.38		129	11%
	17/06/2021– 24/08/2021	< 1.4		< 1.3		141	14%
	24/08/2021– 27/10/2021	< 0.38		< 0.32		139	10%

Table 19. Activity concentrations of the plutonium isotopes Pu-238, Pu-239 and Pu-240 of the bladder wrack samples and sinking matter samples collected from the marine environment of Loviisa and Olkiluoto in 2021.

Site	Site	Sample type	Collection period	Pu-238 Bq/kg	2 σ	Pu-239/240 Bq/kg	2 σ
Loviisa	Stenöarna	Bladder wrack	11/05/2021	< 0.008	40%	0.076	21%
	Hästhölm	Bladder wrack	01/09/2021	0.036		0.053	28%
	Storskarven E	Bladder wrack	12/05/2021	< 0.017		0.039	28%
	Storskarven E	Bladder wrack	31/08/2021	< 0.018		0.044	31%
Olkiluoto	Kalliopöytä	Bladder wrack	05/05/2021	< 0.016		0.043	27%
	Iso Kaalonperä 9	Bladder wrack	12/08/2021	< 0.017		0.043	33%
	Viikari 16	Bladder wrack	04/05/2021	< 0.006		0.030	25%
Loviisa	Hästhölmfjärden 5S	Sinking matter	12/11/2020–18/11/2021	< 0.024	50%	0.90	14%
	Päsalöfjärden R1	Sinking matter	11/11/2020–16/11/2021	0.029		0.31	15%
Olkiluoto	Rääpinkivet 3	Sinking matter	17/11/2020–26/10/2021	< 0.019		0.60	14%
	Keskivedenkari 18	Sinking matter	18/11/2020–27/10/2021	< 0.028		0.47	15%

Radionuclides originating from the power plants were found in surface sediment in the marine environment of the power plants (Table 20). Co-60 and Ag-110m were detected in Loviisa, and the Cs-137 concentration of sediment in the vicinity of the power plants was between 280 and 340 Bq/kg. Co-60 was detected at Olkiluoto, and the Cs-137 activity concentration of sediment in the vicinity of the power plants was between 180 and 200 Bq/kg. The sediment reference samples were collected further away from the power plants and no radionuclides originating from the power plants were detected in these samples, and the Cs-137 activity concentration of the reference samples was 210 – 230 Bq/kg.

Table 20. Radioactive substances found in the marine environment sediment samples in 2021.

Site	Collection date	Co-60 Bq/kg	2 σ	Ag-110m Bq/kg	2 σ	Cs-137 Bq/kg	2 σ
Hästhölmfjärden 5, Loviisa	29/09/2021	0.50	26%	1.22	29%	281	8.6%
Hästhölmfjärden 3, Loviisa	29/09/2021	0.70	53%	0.80	32%	280	11%
Klobbfjärden 1, Loviisa	29/09/2021	< 0.67		< 0.88		338	16%
Hudöfjärden 10, Loviisa	29/09/2021	< 1.0		< 1,1		279	8.1%
Päsalöfjärden R1, Loviisa	29/09/2021	< 0.47		< 0.73		231	16%
Vähä Kivikkokari 12, Olkiluoto	20/05/2021	0.95	27%	< 0.59		197	8.1%
Olkiluoto 9, Olkiluoto	19/05/2021	2.1	18%	< 0.79		200	8.1%
Liponluoto 2, Olkiluoto	19/05/2021	1.3	33%	< 0.66		184	6.6%
Tankarit 4, Olkiluoto	20/05/2021	0.58	39%	< 0.46		179	10%
Olkiluoto S8, Olkiluoto	20/05/2021	< 0.41		< 0.63		213	11%

6.4 Inhabitants of the surroundings

No radioactive substances originating from the power plants were detected in the measurements of inhabitants of the area surrounding the power plants. Measurements were made on 15 individuals in Loviisa and 22 in Olkiluoto.

7 Summary and conclusions

In 2021, small quantities of radioactive substances originating from the power plants were found in the samples collected from the marine environment of both Finnish nuclear power plants. The terrestrial environment samples showed mainly fallout originating from the Chernobyl disaster. The quantities of the radioactive substances correspond largely to those observed in the environment of the plants in the previous years and follow the longer-term downward trend, which is influenced by the development in the control of power plant releases. The quantities of radioactive substances detected in the environment were so small that they are insignificant in terms of the radiation exposure of the environment or people. No radioactive substances originating from the power plants were detected in the measurements of inhabitants of the area surrounding the power plants. The calculated radiation dose of the most exposed individual in the vicinity of both the Loviisa and Olkiluoto nuclear facilities was in 2021 less than 1% of the limit of 0.1 millisieverts set in the Nuclear Energy Decree (161/1988) (Häikiö, 2022).

The Cs-137 concentration observed in the particle samples collected from external air during the annual outages is equivalent to the concentration of Cs-137 found in samples collected elsewhere in Finland, originating mainly from the fallout of the Chernobyl disaster. Similarly, the quantities of Cs-137 and Sr-90 found in the terrestrial environment samples do not differ significantly from corresponding samples collected elsewhere in Finland, and the differences in the concentrations of Cs-137 in the different samples between the plant sites can be explained by the regional differences in the fallout of the Chernobyl disaster across Finland. Moreover, the quantities of Cs-137 and Sr-90 found in the samples do not differ from those found in the terrestrial environment samples of the vicinity of the power plants in the previous years. The C-14 concentrations of samples collected in the surroundings of the plants correspond to the concentrations of the reference samples collected elsewhere in Finland. No other radionuclides, possibly originating from the power plants, were detected in the terrestrial environment samples in 2021.

The Cs-137 concentrations observed in the terrestrial environment samples examined by the licensees correspond to those observed by STUK as part of its own monitoring. In the environment of both power plants, no radionuclides, other than Cs-137, possibly originating from the power plants were detected in the terrestrial environment samples examined by the licensees. The Cs-137 concentration observed in the samples is equivalent to the concentrations of Cs-137 found in environmental samples elsewhere in Finland, originating mainly from the fallout of the Chernobyl disaster.

In the vicinity of both power plants, small amounts of radionuclides originating from the power plants can be seen in the marine environment samples. The radionuclides observed do not fundamentally deviate from the nuclides, originating from the power plants, observed in the marine environment in the previous years. Same nuclides were found in the marine environment of the nuclear facilities as were reported by the power plants to have been released into the environment. The findings of the environmental monitoring of the nuclear facilities carried out by STUK correspond to the findings of the environmental monitoring carried out by the licensees.

8 References

Fortum Power and Heat Oy (2022), Ympäristön säteilyturvallisuuden vuosiraportti 2021.

HELCOM (2018), Thematic assessment of the radioactive substances in the Baltic Sea, 2011 – 2015. Baltic Sea Environment Proceedings No. 151.

IAEA (2005), Environmental and source monitoring for purposes of radiation protection: safety guide, IAEA Safety series standards series No. RS-G-1.8, International Atomic Energy Agency, Vienna, Austria.

Kostiainen, E. and Ylipieti, J. (2010), Radioaktiivinen cesium Suomen ruokasienissä, STUK-A 240. Helsinki, Radiation and Nuclear Safety Authority.

Mattila A. and Inkinen S. (ed.) (2021), Environmental Radiation Monitoring in Finland: Annual Report 2020, STUK-B 268. Helsinki, Radiation and Nuclear Safety Authority.

Siiskonen, T. (ed.) (2020), Suomalaisten keskimääräinen efektiivinen annos vuonna 2018, STUK-A 263. Helsinki, Radiation and Nuclear Safety Authority.

TVO Teollisuuden Voima Oyj (2022), Olkiluodon ydinvoimalaitoksen ympäristön säteilyturvallisuuden vuosiraportti 2021.

Häikiö, J. (ed.) (2022), Regulatory oversight of safety in the use of nuclear energy: Annual Report 2021, STUK-B 280. Helsinki, Radiation and Nuclear Safety Authority.

9 Annexes

- Annex 1** Most commonly found radionuclides in the radiation monitoring of the environment of the nuclear power plants
- Annex 2** Minimum requirements for a nuclear power plant's programme of environmental radiation surveillance, implemented by the licensee
- Annex 3** Schedule for the acquisition of STUK's monitoring samples
- Annex 4** Results of periphyton sample monitoring measurements
- Annex 5** Terrestrial environment results 2020

ANNEX 1

The radionuclides most commonly detected in the environment of nuclear power plants

Nuclide	Half-life	Most common source in environmental samples	Occurrence in environmental monitoring
H-3 tritium	12.2 years	Plant emissions and 1950s and 1960s nuclear weapons tests	Water samples (land and marine environment)
Be-7 beryllium	53 days	Generated in the stratosphere as a result of cosmic radiation	
C-14 carbon	5,700 years	Cosmic (occurring in nature) or from power plants	C-14 from a power plant in gaseous form (CO ₂ or CH ₄), may end up in plants through photosynthesis (if CO ₂ emission).
K-40 potassium	1.248×10 ⁹ years	Naturally occurring radioactive substance	
Cr-51 chrome	27.7 years	Power plant releases	Atmospheric and maritime environment
Mn-54 manganese	312 days	Power plant releases	Atmospheric and maritime environment
Co-58, Co-60, cobalt	70 days 5.3 years	Power plant releases	Atmospheric and maritime environment
Sr-89, Sr-90 strontium	51 days 28.8 years	Power plant releases. Sr-90 in environmental samples also from nuclear weapon testing in the 1950s and 1960s	In maritime and land environment
Ru-103 Ru-106 ruthenium	39 days 372 days	Releases from a power plant or other nuclear facility	In air samples
Ag-110m silver	250 days	Power plant releases	Atmospheric and maritime environment
Sb-124 antimony	60 days	Power plant releases	Atmospheric and maritime environment
I-131 iodine	8 days	Power plant releases, also in use in nuclear medicine at hospitals	May sometimes be detected in the monitoring of air and maritime environment samples, also separately inspected in milk (never discovered). Also detected in sludge samples of water treatment plants, where iodine ends up mainly as a result of medicinal use.
Cs-134, Cs-137 caesium	2.1 a 30 a	Cs-137 in environmental samples mainly from the Chernobyl fallout,	Land and maritime environment

ANNEX 1
 THE RADIONUCLIDES MOST COMMONLY DETECTED IN THE ENVIRONMENT OF NUCLEAR POWER
 PLANTS

		Cs-134 a shorter-lived fission product and from power plant releases	
Ce-141 Ce-144 cerium	33 days 284 days	Power plant releases	Atmospheric and maritime environment
Pu-238, Pu-239 Pu-240 plutonium	87.7 years 24,110 years 6,561 years	Small concentrations detected in environmental monitoring, from nuclear weapon testing in the 1950s and the 1960s	In sediments and in sinking matter


ANNEX 2

Minimum requirements for a nuclear power plant's programme of environmental radiation surveillance, implemented by the licensee (Guide YVL C.7)


Control target	Number of monitoring instruments or samples and measurement or sampling sites	Collection frequency (number/period)	Analysis and frequency
B01. External radiation	External radiation dose rate measuring stations in the site area (or its vicinity) and outside of it at a distance of approx. 5 km from the power plant	—	Continuous measurement and recording
B02. External radiation	10–20 dosimeter stations evenly spread in the key directions at 1–10 km from the power plant	Continuous collection; dosimeters replaced four times a year	Gamma dose 4 times a year
B03. Radioactive substances in the form of airborne particles and iodine in the air	4–5 air sample collectors 1–10 km from the power plant	Continuous collection; filters replaced twice a month, except from the closest collector once a week during annual maintenance	Gamma emitters twice a month (once a week)
B04. Deposition	3–5 rainwater collectors 1–10 km from the plant	Continuous collection; replacement from the closest collector once a month and from the others four times a year	Gamma emitters and ³ H from the closest collector once a month; other gamma emitters and ³ H four times a year.
B05. Indicator organisms in the terrestrial environment	A minimum of one indicator species that enriches radionuclides	1–2 times a year	Gamma emitters 1–2 times a year
B06. Garden products	1–10 km from the power plant; a minimum of 1 species	1–2 times a year	Gamma emitters 1–2 times a year
B07. Domestic water	From the power plant	4 times a year	Gamma emitters and ³ H 4 times per year
B08. Seawater or lake water depending on plant site	From at least one location near the discharge opening	2–4 times a year	Gamma emitters and ³ H from the closest point 4 times a year
B09. Special areas	If necessary, special areas in the environment of the nuclear power plant that may be significant in terms of radiation exposure to the environment, living populations or humans may be selected as control targets. Special areas may include, for example, landfill runoff from the site area, water from the wastewater treatment plant and products grown or farmed near the nuclear facility (such as when residual heat from a power plant is utilized in the production of foodstuffs).		

ANNEX 3

Collection schedule of STUK's monitoring samples



ANNEX 3
COLLECTION SCHEDULE OF STUK'S MONITORING SAMPLES



Collection schedule plan of STUK's monitoring samples.

ANNEX 4

Results of periphyton sample monitoring measurements

Table 21. Results of Loviisa periphyton sample monitoring measurements in 2021.

Collection period	Be-7	2σ	K-40	2σ	Mn-54	2σ	Co-58	2σ	Co-60	2σ	Nb-95	2σ	Zr-95	2σ	Ag-110m	2σ	Sb-124	2σ	Cs-137	2σ
12/11/2020– 22/04/2021	567	17%	392	18%	< 0.84		< 1.2		2.9	23.00 %	< 2.3		< 2.2		2.0	21%	< 1.4		90	11%
22/04/2021– 12/05/2021	457	9.4 %	414	14%	< 1,1		< 1.2		< 1.5		< 1.6		< 2.2		1.7	53%	< 1.3		104	9%
12/05/2021– 27/05/2021	630	12%	518	20%	< 1.4		< 1.5		< 1.8		< 1.9		< 2.6		< 2.2		< 1.7		90	11%
27/05/2021– 01/07/2021	284	10%	651	15%	< 0.38		< 0.40		< 0.44		< 0.56		< 0.77		< 0.48		< 0.42		37	10%
01/07/2021– 03/08/2021	260	12%	402	16%	< 1.0		< 1,1		< 1,1		< 1.6		< 2.1		< 1.3		< 1.2		37	16%
03/08/2021– 19/08/2021	1313	12%	496	18%	< 0.56		< 0.50		< 0.61		< 0.65		< 0.95		< 0.73		< 0.54		107	13%
19/08/2021– 09/09/2021	525	11%	545	16%	1.3	33%	1.1	20%	1.3	23%	21	12 %	13.5	13%	13	16%	1.2	31%	139	11%
09/09/2021– 28/09/2021	393	10%	453	15%	2.6	26%	1.3	44%	7.9	16%	3.9	19 %	2.8	16%	8.0	14%	1.7	34%	87	14%
28/09/2021– 18/11/2021	721	10%	437	15%	< 1.6		< 2.1		< 2.0		< 2.5		< 3.5		1.9	33%	< 2.3		103	9%

ANNEX 4
RESULTS OF PERIPHYTON SAMPLE MONITORING MEASUREMENTS

Table 22. Results of Olkiluoto periphyton sample monitoring measurements in 2021.

Collection period	Be-7	2 σ	K-40	2 σ	Mn-54	2 σ	Co-60	2 σ	Cs-137	2 σ
15/04/2021– 20/05/2021	244	19%	290	21%	< 1.9		< 2.4		11	21%
20/05/2021– 03/06/2021	131	14%	351	17%	< 1.5		< 1.8		25	17%
03/06/2021– 29/06/2021	113	14%	827	13%	< 0.42		1.1	32%	51	18%
29/06/2021– 28/07/2021	35	17%	749	16%	0.71	25%	1.4	29%	20	16%
28/07/2021– 12/08/2021	253	11%	458	16%	< 0.82		< 1.0		23	11%
12/08/2021– 17/09/2021	220	11%	612	13%	< 0.79		1.1	31%	65	14%
17/09/2021– 05/10/2021	533	9%	465	13%	< 0.67		4.5	17%	60	8.8%
05/10/2021– 04/11/2021	537	13%	499	19%	< 0.75		1.5	27%	66	18%

ANNEX 5

Monitoring measurement results of soil samples in 2020

Table 23. Results of Loviisa soil samples (surface layer 0 – 5 cm) in 2020.

Site	Date	K-40 Bq/kg	2σ	Co-60 Bq/kg	2σ	Sr-90 Bq/kg	2σ	Ag-110m Bq/kg	2σ	Cs-137 Bq/kg	2σ
Ingersholm	07/04/2020	880	14	< 0.1		1.5	9	< 0.1		39	10
Bölsängen	08/04/2020	710	16	< 0.1		2.6	10	< 0.2		240	10
Tallholmen	07/04/2020	74	15	< 0.2		5.2	9	< 0.3		1,400	10
Tallholmen 2	08/04/2020	610	12	< 0.1		5.5	9	< 0.2		270	8
Hudö G	14/05/2020	610	12	< 0.2		1.8	9	< 0.3		210	8

Table 24. Results of Olkiluoto soil samples (surface layer 0 – 5 cm) in 2020.

Site	Date	K-40 Bq/kg	2σ	Co-60 Bq/kg	2σ	Sr-90 Bq/kg	2σ	Ag-110m Bq/kg	2σ
Itäranta	15/04/2020	450	15	< 0.2		2.1	9	150	10
Munakari	15/04/2020	670	16	< 0.2		1.9	9	40	10
Flutanperä 2	15/04/2020	89	13	< 0.2		6.5	9	100	8
Viinitarha	16/04/2020	550	15	< 0.2		2.1	9	150	10
Kuusisenmaa	15/04/2020	510	14	< 0.1		2.5	9	71	10